



Climate Change, Variability and Drought Portfolio

Laupāhoehoe Natural Area Reserve

Authors

Ryan J. Longman, East-West Center

Abby G. Frazier, East-West Center

Christian P. Giardina, USDA Forest Service

Hawai'i Drought Knowledge Exchange

Climate change, climate variability, and drought (CCVD) will exert a growing impact on Hawai'i's landscapes, watersheds, and nearshore areas in the future. While managers are tasked with utilizing the best available science, they are often unaware of what datasets or products are available as there is no centralized, drought-focused information clearinghouse or mechanism to engage with scientists in research, planning, product development, and knowledge co-production. The Hawai'i Drought Knowledge Exchange (HDKE) funded by the Pacific Islands Climate Adaptation Science Center, focuses on facilitating knowledge exchange between the research community and resource managers and user groups, thereby expanding the utility of drought-related products for resource managers.



CCVD Portfolio

The climate change, climate variability, and drought (CCVD) portfolio is a comprehensive synthesis of climate and drought information developed specifically for Laupāhoehoe Natural Area Reserve (Laupāhoehoe NAR). This portfolio is designed to provide both research scientists and land managers with relevant climate and drought information needed to inform land management and guide future research and extension.

Laupāhoehoe NAR (Hawaii Island)




Figure 1. Map of Laupāhoehoe NAR.

Part 1: Climate Characteristics

Laupāhoehoe NAR

In developing this Portfolio, we relied on several gridded climate products available for the State of Hawai'i. Both mean annual and mean monthly estimates of rainfall were obtained from the Rainfall Atlas of Hawai'i (<http://rainfall.geography.Hawaii.edu/>). Gridded estimates of other climate variables are obtained from the Climate of Hawai'i (<http://climate.geography.Hawai'i.edu/>). We retrieved all the data points that fell within the boundaries of Laupāhoehoe NAR from our 250 m resolution state-wide maps to support the presented analyses.



Elevation

Laupāhoehoe NAR is located on the Island of Hawai'i Island and covers a vertical elevation range of 4537 ft. In Hawai'i, climate gradients can change significantly over short distances due to changes in elevation, topography, and orientation to the prevailing winds.

Elevation Laupāhoehoe NAR

Minimum = 197 ft

Mean = 3284 ft

Maximum = 4734 ft




Figure 2. Elevation for the Island of Hawai'i Island with Laupāhoehoe NAR outlined in black. The maps shown in the following slides will be for the Laupāhoehoe NAR area only.

Average Annual Climate Characteristics

RF- Rainfall

Spatial Range: 121 to 197 in

RH- Relative humidity

Spatial Range: 82 to 86 %

SM- Soil moisture

Spatial Range: 0.65 to 0.74 Ratio

Mean TA- Average temperature

Spatial Range: 56.4 to 66.3 °F

Min TA- Average minimum temperature

Spatial Range: 48.9 to 60 °F

Max TA- Average maximum temperature

Spatial Range: 65.3 to 74 °F

CF- Cloud cover

Spatial Range: 0.58 to 0.58 Ratio

S- Solar radiation

Spatial Range: 191 to 219 W/m²

ET- Evapotranspiration

Spatial Range: 32 to 49 in

Average Annual Climate: Laupāhoehoe NAR




Figure 3. Mean annual climate of Laupāhoehoe NAR with area average shown in heading of each plot.

Average Monthly Rainfall and Temperature

Average monthly rainfall and temperature patterns vary over the course of the year at Laupāhoehoe NAR. The highest monthly rainfall is received in March (20 in) and the lowest monthly rainfall is received in June (9 in). There is a 6°F annual variation in temperature with the warmest month occurring in September (64 °F) and the coolest month occurring in February (58 °F).

Monthly Climate: Laupāhoehoe NAR





Figure 4. Mean monthly rainfall and temperature at Laupāhoehoe NAR.

Average Monthly Climate

Monthly Rainfall: Laupahoehoe NAR



Monthly Temperature: Laupahoehoe NAR




Figure 5. Mean monthly rainfall Laupāhoehoe NAR with area average shown in heading of each plot.


Figure 6. Mean monthly temperature at Laupāhoehoe NAR with area average shown in heading of each plot.

Average Seasonal Rainfall

Hawai'i has two distinct 6-month seasons of rainfall. The Dry season runs from October to May and the Wet season runs from November to April. At Laupāhoehoe NAR, dry season rainfall averages 11 inches per month (66 inches total) and wet season rainfall averages 16.2 inches per month (97 inches total). Across the entire unit, rainfall ranges from 7.5 to 13.9 inches per month (45 to 84 total) during the dry season and 12.7 to 19 during the wet season (76 to 114 total).

Seasonal Rainfall: Laupāhoehoe NAR

Dry Season (MAY-OCT) 66.2 in



Wet Season (NOV-APR) 97.4 in




Figure 7. Dry and Wet season total rainfall at Laupāhoehoe NAR.

Average Monthly Climate Table

Laupāhoehoe NAR


Variable	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	ANN
RF [in]	14.0	13.0	20.0	18.3	11.1	9.0	13.2	12.8	9.9	10.3	17.1	14.6	164.0
Min TA [°F]	51.5	51.3	51.5	52.5	53.9	55.7	56.3	56.4	56.5	56.1	54.8	52.7	54.1
Mean TA [°F]	58.8	58.4	58.3	59.1	60.7	62.9	63.0	63.1	63.9	63.6	61.2	59.0	61.0
Max TA [°F]	68.0	67.2	66.7	66.8	68.4	70.9	70.6	70.9	72.8	71.9	69.0	66.8	69.2
RH [%]	82.0	82.0	85.0	88.0	87.0	89.0	87.0	86.0	87.0	86.0	84.0	83.0	85.0
CF [%]	48.0	59.0	64.0	63.0	54.0	51.0	47.0	48.0	46.0	52.0	53.0	53.0	53.0
ET [in]	2.0	2.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	3.0	2.0	2.0	33.0
SM [%]	71.0	71.0	74.0	77.0	72.0	65.0	67.0	69.0	67.0	65.0	70.0	73.0	70.0
S [W m/2]	170.0	182.0	194.0	210.0	239.0	236.0	245.0	226.0	214.0	191.0	164.0	154.0	202.0

Table 1. Average monthly climate variables characteristics at Laupāhoehoe Natural Area Reserve. Where, RF is rainfall; Min TA is average minimum air temperature Mean TA is average air temperature; Max TA is average maximum air temperature; RH is relative humidity; CF is cloud frequency; ET is evapotranspiration; SM is soil moisture; S is shortwave downward radiation; ANN, is annual total for rainfall and annual average for all other variables.

Part 2: Inter-Annual Rainfall

Laupāhoehoe NAR

Rainfall in Hawai'i can vary greatly from year-to-year due to natural modes of climate variability such as the El Niño-Southern Oscillation (ENSO). ENSO can be explained as an interaction between the atmosphere and the ocean in the tropical Pacific that results in a somewhat periodic variation between below-normal and above-normal sea surface temperatures. In Hawai'i, wet season rainfall is typically low during the warm (El Niño) phase of ENSO and high during the neutral and Cool (La Niña) Phases. This pattern is reversed in the dry season although it is not as pronounced as in the wet season. ENSO is the dominant mode of climate variability in Hawai'i.



Wet Season Rainfall vs El Niño/La Niña

93-years of monthly wet season rainfall (1920-2012) are compared with the Multivariate ENSO Index (MEI) to determine how rainfall is influenced by five different ENSO phases. During the strong El Niño phase, average monthly wet season rainfall (12 in/month) is 74% drier than the long-term average (16.2 in/month).

Table 2. ENSO phases and abbreviations corresponding to Figure 8.

SEL	Strong El Niño
WEL	Weak El Niño
NUT	Neutral
WLA	Weak La Niña
SLA	Strong La Niña




Figure 8. Boxplot of wet season monthly rainfall grouped by ENSO category. Laupāhoehoe NAR.

Long-Term Trends in Rainfall

Linear trends in annual, wet season, and dry season rainfall are calculated over a 100-year record at Laupāhoehoe NAR for six different periods in the record. Each trend period has a unique start year, but they all end in 2019. When the p-value is less than 0.05, the trend is determined to be statistically significant.




Figure 9. 100-year (1920-2019) rainfall time series at Laupāhoehoe NAR with linear trends calculated over six unique periods in the record; Trend is the slope (inches per year); R2 is the strength of the trend; p is a measure of the statistical significance.


Part 3: Drought and Fire History

Laupāhoehoe NAR

Drought is a prominent feature of the climate system in Hawai'i and can cause severe impacts across multiple sectors. Droughts in Hawai'i often result in reduced crop yields, loss of livestock, drying of streams and reservoirs, depletion of groundwater, and increased wildland fire activity. These impacts can cause substantial economic losses as well as long-term damage to terrestrial and aquatic habitats.



5-Types of Drought



Meteorological Drought.

Defined as a lack of rainfall



Agriculture Drought

Refers to a period of declining soil moisture and subsequent crop failure



Hydrological Drought

Expressed as decreased streamflow and sub-surface water storage



Ecological Drought

Includes any impacts to ecosystems including an increase in wildfire occurrence



Socio-Economic Drought

Includes impacts to social and economic systems, including increased costs or revenue losses, and impacts on public health and safety

Identifying Droughts Using the Standard Precipitation Index

The Standardized Precipitation Index (SPI) is one of the most widely used drought indices. SPI compares rainfall with its multi-year average, and because droughts are generally defined relative to the local normal, this standardized index allows wet and dry climates to be represented on a common scale. Here, Here, 100-years of monthly rainfall are used to used to calculate SPI-12, which compares how a 12-month period compares with all 12-month periods in the record. SPI-12 is a good measure of sustained droughts that affect hydrological processes at Laupāhoehoe NAR.




Figure 10. 100-year (1920-2019) SPI-12 time series at Laupāhoehoe NAR positive SPI (blue) indicate wet periods, negative SPI (red) indicate dry periods.

A 100-Year History of Drought

Negative SPI values (dry periods) are inverted to show a complete drought timeseries at Laupāhoehoe NAR. Dashed lines and corresponding color coding indicates instances of Moderate (SPI > 1), Severe (SPI > 1.5), and Extreme (SPI > 2) drought.

A total of 21 Droughts were observed over the 100-year record with total of 14 drought events of severe strength or greater. The longest drought lasted for a total of 60 months (see Table 2 on next slide).

SPI-12 Drought Events 1920 -2019: Laupahoehoe NAR




Figure 11. 100-year (1920-2019) SPI time series (reversed axis) at Laupāhoehoe NAR. Dashed lines show, moderate (yellow), severe (red), and extreme (dark red), drought thresholds.

Drought Events (1920-2019)

Laupāhoehoe NAR

Start Date	End Date	Duration (months)	Average Intensity	Peak Intensity	Magnitude
1920-12	1921-11	11	0.51	1.87	5.57
1924-08	1927-12	31	1.04	2.57	32.13
1928-01	1935-02	53	0.68	2.29	35.81
1935-12	1936-08	8	0.49	1.12	3.92
1940-01	1941-05	16	1.06	2.15	16.91
1943-03	1947-10	55	0.84	1.79	46.06
1947-12	1950-05	16	0.56	1.53	9.04
1951-04	1954-12	36	0.99	2.12	35.60
1955-12	1964-02	60	0.79	3.14	47.46
1964-07	1967-04	22	0.46	1.05	10.17
1967-10	1975-02	52	0.66	1.30	34.22
1975-04	1976-11	19	0.60	1.68	11.38
1977-01	1979-01	19	0.46	1.11	8.70
1980-01	1982-03	14	1.42	2.43	19.82
1983-07	1985-09	26	0.83	2.00	21.51
1986-02	1993-07	22	0.66	1.49	14.54
1993-08	1997-08	27	0.58	1.15	15.63
1998-02	2004-03	35	0.61	1.32	21.41
2005-04	2009-10	36	0.71	1.72	25.44

Table 2. SPI-12 drought characteristics at Laupāhoehoe Natural Area Reserve identified in the SPI-12 timeseries. Duration is the number of months the drought persisted; Average Intensity is the average absolute SPI; Peak Intensity is the highest SPI value calculated during the drought Magnitude is sum of absolute SPI values during the drought.

Short-term vs Long-term Droughts

SPI-3 Drought Events 1990-2019: Laupāhoehoe NAR




Figure 12. 30-year (1990-2019) SPI-3 time series (reversed axis) at Laupāhoehoe Natural Area Reserve.

SPI-12 Drought Events 1990-2019: Laupāhoehoe NAR




Figure 13. 30-year (1990-2019) SPI-12 time series (reversed axis) at Laupāhoehoe Natural Area Reserve.

The SPI-3 provides a comparison of the precipitation over a specific 3-month period and reflects short- and medium-term moisture conditions over the 30-year period (1990-2019). A total of 16 droughts were observed at Laupāhoehoe NAR. Over this same time period, only 6 droughts were identified when looking at the SPI-12 timeseries. It is important to compare the 3-month SPI with longer time scales. A relatively normal 3-month period could occur in the middle of a longer-term drought that would only be visible at longer time scales. Looking at longer time scales would prevent a misinterpretation that a drought might be over.

Fire Occurrence Hawai'i Island

Ecological drought in Hawai'i is often drives an increase in wildfire occurrence. In Hawai'i, wildfires are most extensive in dry and mesic non-native grasslands and shrublands. During drought events, wildfire risk in grasslands increases rapidly. Changes in land use that shift agricultural land to non-native cover of fire-prone grasses and shrubs combined with recurring incidences of drought are expected to increase the risk of future wildfire in Hawai'i.





Figure 14. The map shows wildfires that have occurred on the island of Hawai'i Island between 1999 and 2019.

Part 4: Future Climate

Laupāhoehoe NAR

To simulate future rainfall and temperature, Global Climate Models are used. These can simulate future conditions under different scenarios for how much carbon dioxide we emit into the air. Two common scenarios are used: RCP 4.5 which assumes we reduce our carbon emissions, and RCP 8.5, a high emissions scenario. The outputs from global models are too coarse to accurately capture changes over the complex terrain of Hawai'i. Therefore, we use an additional step called Downscaling to relate the global-scale information down to the local island scale.

In Hawai'i, two types of downscaled projections are available.
Dynamical Downscaling (End of Century)
Statistical Downscaling (Mid & End of Century)
Results for both types of downscaling and both scenarios will be shown here.



Average Rainfall Change 2100 Year 2100

Rainfall in the Year 2100

Annual

2 to 7 in/year
(1 to 4 % Change)

Dry Season

-1 to 1 in/month
(-7 to 5 % Change)


Wet Season

-1 to 2 in/month
(-4 to 12 % Change)


The range in projections include estimates for both low emissions (RCP 4.5) and high emissions (RCP 8.5) scenarios, and for both Dynamical and Statistical Downscaling approaches.

Dynamical and Statistical DS RCP 8.5, 2100


DyDs ANN (2%)




StDs ANN (4%)




DyDs Dry (-7%)



StDs Dry (-6%)



DyDs Wet (-4%)



StDs Wet (12%)




Figure 15. Downscaled future rainfall projections (% Change; (2100) at Laupāhoehoe NAR, Dynamical Downscaling (DyDs), Statistical Downscaling (StDs), for annual (ANN), dry season (DRY) and wet season (WET).

Average Rainfall Change 2040-2070

Rainfall for Years 2040-2070

Annual

2 to 7 in/year
(1 to 4% Change)

Dry Season

0 to 0 in/month
(-3 to -3% Change)


Wet Season

1 to 1 in/month
(3.1 to 8% Change)


The range in projections include estimates for both low emissions (RCP 4.5) and high emissions (RCP 8.5) scenarios, for the Statistical Downscaling approach.

Statistical DS RCP 8.5 & RCP 4.5, 2040-2070)


StDs ANN (1%)




StDs ANN (4%)




StDs DRY (-3%)



StDs DRY (-3%)



StDs WET (4%)



StDs WET (8%)




Figure 16. Downscaled future rainfall projections (% Change; 2040-2070) at Laupāhoehoe NAR, for the Statistical Downscaling (StDs) approach, for annual (ANN), dry season (DRY), and wet season (WET) for RCP 4.5 (left) and RCP 8.5 (right).

End-of-Century Change in Temperature

Mean Temperature Now

61°F

Mean Temperature 2100


64.1°F to 66.9°F

(3.1°F to 5.9°F Change)


The range in projections include estimates for both low emission (RCP4.5) and high emissions (RCP8.5) scenarios, for both the Dynamical and Statistical Downscaling approaches.

Dynamical & Statistical DS Compare TEMP (Yr 2100)


DyDs ANN RCP 4.5 (3.2 °F)



StDs ANN RCP 4.5 (3.1 °F)



DyDs ANN RCP 8.5 (3.2 °F)



StDs ANN RCP 8.5 (5.9 °F)




Figure 17. Downscaled projected change in mean temperature (Year 2100) at Laupāhoehoe NAR, Dynamical Downscaling (DyDs), Statistical Downscaling (StDs).

Mid-Century Change in Temperature

Mean Temperature Now

61°F

Mean Temperature 2040-2070


63.4°F to 64.4°F

(2.4°F to 3.4°F Change)

The range in projections include estimates for both low emission (RC 4.5) and high emissions (RCP 8.5) scenarios, for the Statistical Downscaling approach.

Statistical DS Compare TEMP (Yr 2040-2070)

StDs ANN RCP 4.5 (2.4 °F)



StDs ANN RCP 4.5 (3.4 °F)




Figure 18. Downscaled Future temperature projections (2040-2070) at Laupāhoehoe NAR for the Statistical Downscaling (StDs) approach.

Part 5: CCVD Summary

Laupāhoehoe NAR

Laupāhoehoe Natural Area Reserve (Laupāhoehoe NAR) is located on the island of Hawai'i Island at mean elevation of 3284 ft (range: 197 to 4734 ft). Rainfall varies over the course of the year with a maximum of 20 inches, occurring in March and a minimum of 9 inches occurring in June. On average, wet season months (Nov-Apr) receive 5.2 in of more rainfall than dry season months (May-Oct). Seasonal rainfall can vary within the unit as well, with wet season rainfall ranging from 45 to 84 inches and wet season rainfall ranging from 76 to 114 inches across the 4537 ft elevation gradient. Rainfall can also vary considerably from year-to-year with the driest years occurring during a Strong El Niño event, when on average, 74% less rainfall is received, relative to the long-term average. The average temperature at Laupāhoehoe NAR is 61°F but temperature ranges from 58°F to 64°F over the course of the year. Drought is a reoccurring feature in the climate system of Laupāhoehoe NAR with a total of 21 occurring over the record which is approximately 2.1 per decade. A total of 14 drought events were at severe strength or greater and the longest drought lasted for a total of 60 consecutive months. Future projections of rainfall are uncertain, with end-of-century annual changes ranging from 1 to 4% and more pronounced changes occurring during the wet and dry seasons. Future projections of temperature suggest an increase of 2.4°F to 3.4°F by mid century (2040-2070) and an increase of 3.1°F to 5.9°F by the end of the century (2100).

External Resources

For more Information

US Drought Monitor

<https://droughtmonitor.unl.edu/>

State of Hawai'i Drought Plan

<https://files.Hawai'i.gov/dlnr/cwrp/planning/HDP2017.pdf>

Rainfall Atlas of Hawai'i

<http://rainfall.geography.Hawai'i.edu/>

Climate Of Hawai'i


<http://climate.geography.Hawai'i.edu/>

ENSO Current Phase and Discussion

https://www.cpc.ncep.noaa.gov/products/analysis_monitoring/enso_advisory/enso_disc.shtml


Pacific Fire Exchange

<https://www.pacificfireexchange.org/>



Acknowledgements

Mari-Vaughn Johnson, Heather Kerkering, Darcy Yogi (PI-CASC), Victoria Keener, Laura Brewington (East-West Center Pacific-RISA), Melissa Kunz, Clay Trauernicht (NREM, UH Manoa), Katie Kamelamela (USDA, Forest Service), Thomas Giambelluca (WRRC, UH Manoa), and John Marra (NOAA).



Suggested Citation

Longman, R.J, Frazier, A.G., Giardina, C.P (2020). Climate Change, Climate Variability and Drought Portfolio Laupāhoehoe Natural Area Reserve Hawai'i Drought knowledge Exchange CCVD Series Volume DRAFT, 31pp.

WARNING: This is a draft. For the most up-to-date version or information about publication please contact Ryan Longman: rlongman@Hawaii.edu

Works Cited

- Elison Timm, O., 2017: Future warming rates over the Hawaiian Islands based on elevation-dependent scaling factors. *Int. J. Climatol.*, **37**, 1093–1104, doi:10.1002/joc.5065.
- , T. W. Giambelluca, and H. F. Diaz, 2015: Statistical downscaling of rainfall changes in Hawai'i based on the CMIP5 global model projections. *J. Geophys. Res.*, **120**, 92–112, doi:10.1002/2014JD022059.
- Frazier, A. ., and Coauthors, A century of spatial and temporal patterns of drought in Hawai'i across hydrological, ecological, and socioeconomic scales. *In Review. Sci. Total Environ.*,.
- Frazier, A. G., T. W. Giambelluca, H. F. Diaz, and H. L. Needham, 2016: Comparison of geostatistical approaches to spatially interpolate month-year rainfall for the Hawaiian Islands. *Int. J. Climatol.*, doi:10.1002/joc.4437.
- Giambelluca, T. W., Q. Chen, A. G. Frazier, J. P. Price, Y. L. Chen, P. S. Chu, J. K. Eischeid, and D. M. Delporte, 2013: Online rainfall atlas of Hawai'i. *Bull. Am. Meteorol. Soc.*, **94**, 212–316, doi:10.1175/BAMS-D-11-00228.1.
- Giambelluca, T. W., and Coauthors, 2014: Evapotranspiration of Hawai'i Final Report.
- Lucas, M. P., R. J. Longman, T. W. Giambelluca, A. F. Frazier, and J. H. Lee, Mapping 30-years of monthly rainfall using an automated kriging approach. *In Review. Data*,.
- Trauernicht, C., 2020: Hawaii Fire Occurrence Map (2000-2019). *Unpublished*,.
- Wolter, K., and M. S. Timlin, 2011: El Niño/Southern Oscillation behaviour since 1871 as diagnosed in an extended multivariate ENSO index (MEI.ext). *Int. J. Climatol.*, **31**, 1074–1087, doi:10.1002/joc.2336.
- Zhang, C., Y. Wang, K. Hamilton, and A. Lauer, 2016: Dynamical downscaling of the climate for the Hawaiian islands. Part II: Projection for the late twenty-first century. *J. Clim.*, **29**, 8333–8354, doi:10.1175/JCLI-D-16-0038.1.

Annex I: 100-Year Rainfall

The 100-year monthly rainfall dataset was drawn from two unique gridded products. We used data from Frazier et al. (2016) for the period 1920-1989 and Lucas et al. (In Review) for the period 1990-2019. Given that two unique data sets and methods were used to make these two products we show the 1:1 Statistical relationship between the two products for a 23-year overlap (1990-2012) with the datasets and associated error metrics.

Frazier et al., 2016; Lucas et al.,
(In Review)




Figure A1: One to one comparison of 23-years (1990-2012) of monthly rainfall from two unique datasets for Laupāhoehoe NAR, and associated error metrics; R2, is the coefficient of determination, MBE, is the mean bias error, MAE, mean absolute error.

Annex II: Climate Downscaling in Hawai'i

Two types of downscaling products were used in this analysis. Here we explain some of the nuances between the two. Dynamical Downscaling (Zhang et al., 2016), feeds GCM output into a regional model that can account for local topographic and atmospheric phenomena at much finer resolutions (e.g. 1 km). End-of-century projections (2100) encompass the period 2080-2099. Statistical Downscaling (Elison Timm et al., 2015, Elison Timm, 2017), develops a relationship between GCM model output and station data for a historical period and then uses this established relationship to make projections for two future scenarios. End-of-century projections (2100) encompass the period 2070-2099 (2100), Mid-century projections encompass the period 2040-2070.

